

**Estimated Total Tons Lost per Year due to Wind Induced Erosion
on Cropland, Conservation Reserve Program Land*, and Pastureland
Ten Counties* in Oregon by NRCS Basin***

National Resources Inventory Results
Estimated for Non-Federal Lands Only

NRCS Basin Name	1982		1987		1992		1997	
	Estimates	Error	Estimates	Error	Estimates	Error	Estimates	Error
--- Tons/Year ---								
John Day/Umatilla	6,058,100	471,400	6,048,300	458,600	4,133,400	397,300	4,134,400	445,300
Deschutes	1,041,500	62,900	1,019,400	59,000	826,500	77,900	899,100	71,500
High Desert	681,800	246,100	632,100	168,000	449,000	172,500	386,200	155,700
Snake River	350,800	86,300	342,700	62,200	288,100	65,000	246,200	75,200
NRCS Basins where wind erosion data not collected due to low levels of wind erosion.								
Central Coast/Upper Willamette	-	-	-	-	-	-	-	-
Lower Willamette	-	-	-	-	-	-	-	-
North Coast	-	-	-	-	-	-	-	-
Southwest Oregon	-	-	-	-	-	-	-	-
Oregon Totals	8,132,200	540,600	8,042,500	496,100	5,697,000	444,800	5,665,900	482,900

* See Table Details

NOTE: Data may not add to totals because of rounding. The error referred to in the table is the standard error of the estimate. You can obtain the margin of error at the 95% confidence limit by multiplying the error by 1.96.

Standard errors greater than 50% of the estimate indicate a low level of statistical reliability.